arXiv:1303.6583v1 [astro-ph.CO] 26 Mar 2013

A Lower Growth Rate from Recent Redshift Space Distortions than Expected from Planck

E. Macaulay[*]
Astrophysics, University of Oxford, DWB, Keble Road, Oxford, OX1 3RH, UK

I. K. Wehus[†]
Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA 91109, USA and
Astrophysics, University of Oxford, DWB, Keble Road, Oxford, OX1 3RH, UK

H. K. Eriksen[‡]
Institute of Theoretical Astrophysics, University of Oslo, P.O. Box 1029 Blindern, N-0315 Oslo, Norway

We perform a meta-study of recently published Redshift Space Distortion (RSD) measurements of the cosmological growth rate, $f(z)\sigma_8(z)$. We analyse the latest results from the 6dFGS, BOSS, LRG, WiggleZ and VIPERS galaxy redshift surveys, and compare the measurements to expectations from Planck. In this Letter we point out that the RSD measurements are consistently less than the values expected from Planck, and the relative scatter between the RSD measurements is lower than expected. Adopting the equation-of-state of dark energy, w, as a strawman for comparison purposes, we obtain a best-fit value of $w = -0.74 \pm 0.07$ from a joint fit to all recent measurements, in tension with the ΛCDM value of $w = -1$ at greater than the three standard deviation level. However, the χ^2 of the resulting fit is 2.70 for 8 degrees of freedom, corresponding to a probability-to-exceed (PTE) of 0.952. On fitting for σ_8, we find σ_8=.722 ± 0.038, χ^2=2.91 and PTE = 0.940. Further, a simple constant in redshift for $f(z)\sigma_8(z)$ provides an even better fit, with $\chi^2 = 2.09$ for a PTE of 0.978. On changing the LRG results to an analysis with a longer maximum correlation scale, we find $w = -0.78 \pm 0.07$, $\chi^2 = 4.01$, and PTE=0.856, a best fit constant value of 0.425 ±0.04, $\chi^2 = 3.71$, and PTE=0.883, or σ_8=.741 ± 0.034, χ^2=4.05 and PTE = 0.852. A full resolution of this issue may require a more robust treatment of non-linear effects in RSD models, although the trend for a low σ_8 agrees with recent constraints on σ_8 and Ω_m from Sunyaev-Zeldovich cluster counts identified in Planck.

Measurements of the growth rate of the cosmological matter density contrast are emerging as an increasingly important cosmological probe, particularly for dark energy. A key advantage of growth rate measurements is that they probe dark energy independently of the background cosmic expansion, so can potentially distinguish between dark energy models which would otherwise be indistinguishable at the background level. An increasingly curious picture may appear to be emerging from several independent surveys, which consistently measure a growth rate lower than the value expected of the ΛCDM model with Planck parameters (Planck Collaboration et al. 2013a). However, the scatter in the collected growth rate measurements is smaller than expected from the measured uncertainties. This suggests that the quoted uncertainties on growth rate measurements may be overestimated, or, alternatively, that some part of the analysis is consistently suppressing scatter in the measurements (or a combination of both cases). We briefly review how the peculiar velocities of galaxies can be used to measure the growth rate, then collect current growth rate measurements, and find a low χ^2 per degree of freedom for a range of w, σ_8, and growth index γ. We then consider possible explanations for these results.

The growth rate f is defined as $f = d\ln G/d\ln a$, where a is the scale factor, and G is the growth factor of the density contrast. Galaxy surveys are directly sensitive to $f(z)\sigma_8(z)$, where σ_8 is the power spectrum normalisation on scales of 8 h^{-1} Mpc. The growth rate is often approximated as $f(z) = \Omega_m(z)^\gamma$, where $\gamma = \frac{6}{11}$ in General Relativity with a cosmological constant (Linder 2005). We use this equation for $f(z)$ only to fit for γ, combined with $\sigma_8(z)$ calculated with fixed Planck parameters. To generate growth rate models for w and σ_8 we use CAMB (Lewis et al. 2000) with HALOFIT (Smith et al. 2003). Since we cannot set σ_8 directly, we vary the amplitude A_s, which is directly related to σ_8.

Most recent growth rate measurements come from inferring peculiar velocities indirectly from Redshift Space Distortions (RSDs) in a galaxy redshift survey, as proposed by Kaiser (1987). This anisotropy was first measured and used to estimate the growth rate by the 2dFGRS (Peacock et al. 2001). The growth rate has since been measured with a range of other techniques and surveys, with the RSD technique in the VVDS survey (Guzzo et al. 2008), QSO clustering and Ly_α clustering (da Ângela et al. 2008; Ross et al. 2007; Viel et al. 2004), and at $z \sim 0$ in peculiar velocity surveys, (Davis et al. 2011; Hudson & Turnbull 2012). In this letter, we con-

---

[*] Electronic address: edward.macaulay@astro.ox.ac.uk
[†] Electronic address: i.k.wehus@fys.uio.no
[‡] Electronic address: h.k.k.eriksen@astro.uio.no

Survey	z	$f(z)\sigma_8(z)$	Reference
6dFGS	0.067	0.423 ± 0.055	Beutler et al. (2012)
LRG_{200}	0.25	0.3512 ± 0.0583	Samushia et al. (2012)
	0.37	0.4602 ± 0.0378	
LRG_{60}	0.25	0.3665 ± 0.0601	
	0.37	0.4031 ± 0.0586	
BOSS	0.30	0.408 ± 0.0552	Tojeiro et al. (2012)
$\rho=-0.19$	0.60	0.433 ± 0.0662	
WiggleZ	0.44	0.413 ± 0.080	Blake et al. (2012)
$\rho=0.51$	0.60	0.390 ± 0.063	
$\rho=0.56$	0.73	0.437 ± 0.072	
VIPERS	0.8	0.47 ± 0.08	de la Torre et al. (2013)

TABLE I: Compilation of recent published values of $f(z)\sigma_8(z)$ (ordered by redshift). We indicate the correlation coefficient ρ where the measurements are correlated in adjacent redshift bins (the 1st and 3rd redshift bins in the WiggleZ survey are uncorrelated).

Parameter	ΛCDM, Planck	Best fit	χ^2	PTE
w	-1	-0.74 ± 0.07	2.70	0.952
σ_8	0.834 ± 0.027	0.722 ± 0.038	2.91	0.940
γ	$\frac{6}{11}$	0.778 ± 0.16	3.18	0.923
Constant		0.415 ± 0.042	2.09	0.978
w	-1	-0.78 ± 0.07	4.01	0.856
σ_8	0.834 ± 0.027	0.741 ± 0.034	4.05	0.852
γ	$\frac{6}{11}$	0.740 ± 0.14	4.05	0.853
Constant		0.425 ± 0.038	3.71	0.883

TABLE II: χ^2 and corresponding Probability To Exceed (PTE) for the best fits to w, σ_8, γ and also a single constant value for the growth rate. We indicate the expected values of w and γ in ΛCDM, and the measured value of σ_8 from Planck Collaboration et al. (2013a). The first set of results is for the LRG_{60} data set, and the second set is for LRG_{200}. Both data sets have 8 degrees of freedom.



FIG. 1: Comparing recent measurements of $f(z)\sigma_8(z)$ to models for a range of w. We are plotting results for the LRG_{60} data set. The measurement error bars are at the 1 standard deviation uncertainty level. The dashed line is the expected growth rate from ΛCDM with Planck parameters, and the solid line is the best-fit growth rate for a variable w, with the 1 and 2 standard deviation uncertainty illustrated with the shaded regions. We note that all the measurements include our best fit model at the 1 standard deviation uncertainty level, which is reflected in the low χ^2 in Table II.

sider only growth rate measurements with RSDs, with published values of $f(z)\sigma_8(z)$, as summarised in Table I.

We consider the χ^2 statistic, given by

$$\chi^2 = (x - \bar{x})\, C^{-1} (x - \bar{x}) \qquad (1)$$

where x is a vector of observed values, $\bar{x}$ is a vector of corresponding values from a model for x, and C is the covariance matrix for the measurements.

In Tojeiro et al. (2012), the growth rate from the BOSS survey was fitted at four correlated redshift values, although the publicly available covariance matrix is for three redshift measurements, to reduce correlations between the measurements. We find that even with three redshift bins, the block-diagonal covariance matrix is too highly correlated, and thus we do not include the highly correlated intermediate redshift measurement. We analyse the data with two different measurements from the LRG survey, with a maximum correlation length of 200 h^{-1}Mpc (LRG_{200}) and also with a maximum correlation length of 60 h^{-1}Mpc (LRG_{60}) – we do not analyse the data with both LRG_{200} and LRG_{60} simultaneously. For both data sets, we calculate conditional best fit parameters, and the corresponding χ^2 for w, σ_8, and γ, with other parameters fixed. For comparison, we also fit the measurements to a single, constant value of $f(z)\sigma_8(z)$ that does not vary with redshift. The results are summarised in table II.

We note that in fact the best fit to the data (in terms of minimum χ^2) is a single, constant value, that is, there is no preference for a model with any change in the growth rate. We find when varying w a best fit value > -1, or when varying σ_8 a best fit value less than the Planck value of 0.834 ± 0.027 (Planck Collaboration et al. 2013a). When varying γ, we find a best fit value higher than expected in General Relativity. In Figure 1 we plot the growth rate measurements from Table I, with growth rate models for ΛCDM, and also for our best fit $w = -0.74 \pm 0.07$ with the LRG_{60} data set. We note that the PTE decreases with the LRG_{200} data set, since the LRG_{200} measurements have a larger scatter than the LRG_{60} measurements. This is likely due to the fact that most of the coherent clustering signal is due to correlations on scales less than 100 h^{-1}Mpc, so the additional correlations are effectively adding noise to the signal.

Although we find that at face value, the current growth rate measurements appear to suggest $w > -1$ or a low value of σ_8, we also emphasise the low χ^2 / n_{dof} we find in the data. While only additional observations will determine whether this trend is truly statistically significant, the results already in hand appear to suggest that either the quoted uncertainties have been overestimated, or the analysis is suppressing genuine scatter in the measurements.

In most recent results, the uncertainties have been estimated from several hundred simulated realisations of the survey, from which the uncertainty (and the covariance between measurements, in the case of several redshift bins) can be deduced from the scatter in the realisations. Although it may appear that the uncertainties on the measurements have been overestimated, good agreement between the quoted values and Fisher forecasts (e.g., White et al. 2009) of the minimum intrinsic statistical uncertainties suggests that this is not the case, although Reid et al. (2012) note that the uncertainties in the BOSS growth rate measurements are around 40% larger than the Fisher matrix predictions.

Perhaps the stage of an RSD analysis most likely to introduce a systematic shift, and artificially reduce the scatter, may be in fitting a model to the two-dimensional two-point correlation function (or power spectrum). Parkinson et al. (2012) analysed simulated catalogues for the WiggleZ survey with a range of models for the RSD effect, and found that measurements of Ω_m (which is directly sensitive to the growth rate) were highly dependent on the model used. In particular, the model of a HALOFIT $P(k)$ with a linear model for the redshift space distortion recovered a lower Ω_m compared to the fiducial value on which the simulation was based.

The preference for a lower growth rate or σ_8 appears to agree with recent results from Planck Collaboration et al. (2013b), studying Sunyaev-Zeldovich (SZ) cluster counts, who find $\sigma_8 = 0.77 \pm 0.02$ and $\Omega_m = 0.29 \pm 0.02$. Collectively, these results may be suggesting that ΛCDM does not fully model simultaneously the Cosmic Microwave Background and the Universe at $z < 1$. However, future work will require detailed work with simulated catalogues for a range of cosmological models (e.g., Jennings 2012; Jennings et al. 2012) and an improved understanding of the relationship between the observed galaxies, the peculiar velocity field, and the underlying dark matter (e.g., Reyes et al. 2010; Sherwin et al. 2012), before we can more robustly use RSD measurements to study departures from ΛCDM.

We thank Erminia Calabrese for help with the Planck parameters for the growth rate, Lado Samushia for help with the growth rate models, Chris Blake and Rita Tojeiro for help with the WiggleZ and BOSS covariance matrices, and Pedro Ferreira for useful comments and discussions. This project was supported by an ERC Starting Grant StG2010-257080 and a Leverhulme visiting professorship for HKE. EM acknowledges support from the BIPAC and STFC. IKW acknowledges support from ERC grant 259505. Part of the research was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with NASA.

Beutler, F. et al. 2012, MNRAS, 423, 3430, 1204.4725
Blake, C. et al. 2012, MNRAS, 425, 405, 1204.3674
da Ângela, J. et al. 2008, MNRAS, 383, 565, arXiv:astro-ph/0612401
Davis, M., Nusser, A., Masters, K. L., Springob, C., Huchra, J. P., & Lemson, G. 2011, MNRAS, 413, 2906, 1011.3114
de la Torre, S. et al. 2013, ArXiv e-prints, 1303.2622
Guzzo, L. et al. 2008, Nature (London) , 451, 541, 0802.1944
Hudson, M. J., & Turnbull, S. J. 2012, APJL, 751, L30, 1203.4814
Jennings, E. 2012, MNRAS, L533, 1207.1439
Jennings, E., Baugh, C. M., Li, B., Zhao, G.-B., & Koyama, K. 2012, MNRAS, 425, 2128, 1205.2698
Kaiser, N. 1987, MNRAS, 227, 1
Lewis, A., Challinor, A., & Lasenby, A. 2000, Astrophys. J., 538, 473, astro-ph/9911177
Linder, E. V. 2005, PRD, 72, 043529, arXiv:astro-ph/0507263
Parkinson, D. et al. 2012, ArXiv e-prints, 1210.2130
Peacock, J. A. et al. 2001, Nature, 410, 169, arXiv:astro-ph/0103143
Planck Collaboration et al. 2013a, ArXiv e-prints, 1303.5076
——. 2013b, ArXiv e-prints, 1303.5080
Reid, B. A. et al. 2012, MNRAS, 426, 2719, 1203.6641
Reyes, R., Mandelbaum, R., Seljak, U., Baldauf, T., Gunn, J. E., Lombriser, L., & Smith, R. E. 2010, Nature (London) , 464, 256, 1003.2185
Ross, N. et al. 2007, MNRAS, 381, 573
Samushia, L., Percival, W. J., & Raccanelli, A. 2012, MNRAS, 420, 2102, 1102.1014
Sherwin, B. D. et al. 2012, Phys. Rev. D , 86, 083006, 1207.4543
Smith, R. E. et al. 2003, MNRAS, 341, 1311, arXiv:astro-ph/0207664
Tojeiro, R. et al. 2012, MNRAS, 424, 2339, 1203.6565
Viel, M., Haehnelt, M. G., & Springel, V. 2004, MNRAS, 354, 684, arXiv:astro-ph/0404600
White, M., Song, Y.-S., & Percival, W. J. 2009, MNRAS, 397, 1348, 0810.1518